

Unifying Scalars and Light Mesons by Holographic QCD

W. de Paula and T. Frederico

Departamento de Física, Instituto Tecnológico de Aeronáutica, 12228-900 São José dos Campos, SP, Brazil

We show that the universality of the infrared dynamics of string modes dual to $q\bar{q}$ states within a Dynamical AdS/QCD model of coupled dilaton-gravity background implies in the Regge-like spectrum of f_0 's scalars and higher spin mesons consistent with experimental data. The pion mass and its trajectory were also described with a scale deformation of the metric and a rescaled string mass. The available experimental decay widths of the $S \rightarrow PP$ decays provided a complementary check of the proposed classification scheme for $f_0(980)$, $f_0(1370)$, $f_0(1500)$ and $f_0(1710)$ as radial excitations of $f_0(600)$. For $f_0(980)$ we estimated a mixing angle of $\pm 20^\circ$ with other structures.

PACS numbers: 11.25.Tq, 11.25.Wx, 14.40.-n, 12.40.Yx

The light scalar mesons are still challenging our imagination and models for quark spectroscopy [1]. Nowadays, tetraquarks configurations [2] are considered dominant for the structure of $f_0(600)$ and $f_0(980)$ scalar resonances [3]. Alternatively, we find that these states can still be interpreted as $q\bar{q}$ states in the framework of a Dynamical AdS/QCD model [4]. The model consists in the solution of five-dimensional gravity coupled to a dilaton, where the metric comes from a deformed anti-de Sitter space in order to have confinement by the area law behavior of the Wilson loop [5].

Besides the masses of the f_0 's, the ratio between $\pi\pi$ decay widths ($\Gamma_{\pi\pi}$) of these mesons are consistent with this assignment. An overall normalization constant for the effective operator that gives the two-pion decay amplitude is introduced and no other free parameter is used in the present calculation beyond those implicit contained in the description of the scalar and pion Regge trajectories. We made the assumption that the holographic coordinate in metric of the Dynamical AdS/QCD model for high spin mesons [4] can be rescaled in order to provide the effective potential for the string modes dual to the scalar and pseudoscalar mesons. The masses of the f_0 scalars are then obtained by a single rescale of the holographic coordinate in the metric form. We also describe the light pseudoscalar states. The scale of the holographic coordinate is contracted for the pseudoscalars in respect to the scalar case and a rescaled string mass is also introduced to allow a description of the pion mass.

The outcome from our work is that $f_0(980)$, $f_0(1370)$, $f_0(1710)$, $f_0(2020)$, $f_0(2100)$, $f_0(2200)$ and $f_0(2330)$ are excited states of $f_0(600)$, and they belong to a Regge trajectory with a slope of about 0.5 GeV^2 half of the corresponding one for the ρ -meson trajectory.

Additionally to support our interpretation of the nature of the light f_0 's scalars, we obtain the wave functions of the string modes dual to the scalar meson states and using the pion string amplitude we calculate the decay width of the scalars into two pions. The experimental observation of the systematic decrease found for $\Gamma_{\pi\pi}$ for higher excitations is easily understood within the holographic view: the overlap of the radial excitations and

the two pion state amplitudes, are enhanced or depleted depending on the number of nodes, and from that the decreasing pattern for the $S \rightarrow PP$ decay width appears.

The two-pion partial decay width for $f_0(600)$, $f_0(1370)$, $f_0(1500)$ and $f_0(1710)$, for which experimental information on the two-pion decay width exists are qualitatively consistent with the model results, using one normalization parameter. In particular for $f_0(600)$ a width of about 500 MeV is found. The experimental value of $\Gamma_{\pi\pi}$ for $f_0(1500)$, known experimentally with a few % error, is used for normalization of the effective action for the decay amplitude in our holographic model.

Although $f_0(980)$ has a mass identified with the first excitation of the string mode dual to $q\bar{q}$ state, it has a too large $\Gamma_{\pi\pi}$ compared to the range of the experimental values. It is known that $f_0(980)$ should mix strongly with other structures such as $s\bar{s}$ states (see e.g. [6]). Indeed we got a mixing angle of about $\pm 20^\circ$ [1] to be consistent with the experimental data for the partial decay width. (We do not determine the sign of the mixing angle.) Below we briefly discuss the basis of present holographic model and substantiate quantitatively our claims.

The search for string dual models of Gauge theories was pursued since the pioneer work by 't Hooft [7] and vigorously developed after the Maldacena's Conjecture [8]. Applications of AdS/CFT to describe QCD began with the Hard Wall model [9] that provides a good description of form factors at high Q^2 but does not give the linear Regge trajectory presented by the mesonic data. This phenomenological result can be obtained by a spin dependence in the metric [10] or by introducing a dilaton field [11]. In particular, scalar mesons were analyzed in ([12] and [13]). However, both works do not include the sigma meson in their studies. Those models are not solutions of Einstein equations and also do not confine by the Wilson Loop criteria. We proposed a Dynamical AdS/QCD model [4] that is a solution of the Einstein equations, confines by the Wilson loop criteria and describe the scalar sector. We start from the Einstein-Hilbert action of five-dimensional gravity coupled to a

dilaton field Φ ,

$$S = \frac{1}{2\kappa^2} \int d^4x dz \sqrt{|g|} \left(-R + \frac{1}{2} g^{MN} \partial_M \Phi \partial_N \Phi - V(\Phi) \right), \quad (1)$$

where κ is the five-dimensional Newton constant, the dilaton field $\Phi(z)$ depends on the radial coordinate only and $V(\Phi)$ is the potential for the dilaton field. We restrict our metric to $g_{MN} = e^{-2A(z)} \eta_{MN}$ with $\eta = \text{diag}(1, -1, -1, -1, -1)$

The static solutions of the Einstein-dilaton coupled field equations in the fifth dimension found in [4] are given by $\Phi' = \sqrt{3} \sqrt{A'^2 + A''}$ and $V(\Phi(z)) = 3e^{2A(z)} [A''(z) - 3A'^2(z)]/2$.

The boundary condition on the physical brane restricts the geometry to asymptotically AdS_5 (AAdS_5) spacetimes and thus ensures conformality in the ultraviolet (UV). The infrared (IR) behavior is chosen to obtain Regge-trajectories consistent with the Wilson-loop area-law for the gravity dual of a confining theory. The conformal invariance is broken in the IR by Λ_{QCD} .

To calculate the spectrum of scalar mesons φ we start from the action

$$I = \frac{1}{2} \int d^4x dz \sqrt{|g|} \left(g^{\mu\nu} \partial_\mu \varphi(x, z) \partial_\nu \varphi(x, z) - \frac{M_5^2}{\Lambda_{QCD}^2} \varphi^2 \right), \quad (2)$$

that describes a scalar mode propagating in the dilaton-gravity background. We factorize the holographic coordinate dependence as $\varphi(x, z) = e^{iP_\mu x^\mu} \varphi(z)$, $P_\mu P^\mu = m^2$.

The string modes of the massive scalar field φ can be rewritten in terms of reduced amplitudes $\psi_n = \varphi_n \times e^{-(3A+\Phi)/2}$ which satisfy the Sturm-Liouville equation

$$[-\partial_z^2 + \mathcal{V}(z)] \psi_n = m_n^2 \psi_n \quad (3)$$

where the string-mode potential is

$$\mathcal{V}(z) = \frac{B'^2(z)}{4} - \frac{B''(z)}{2} + M_5^2 e^{-2A(z)}, \quad (4)$$

with $B = 3A + \Phi$. (Note that $B = (2S - 1)A + \Phi$ for the spin nonzero states [11].) The gauge/gravity dictionary identifies the eigenvalues $m_{n,S}^2$ with the squared meson mass spectrum of the boundary gauge theory.

The AdS/CFT correspondence states that the wave function should behave as z^τ , where $\tau = \Delta - \sigma$ (conformal dimension minus spin) is the twist dimension for the corresponding interpolating operator that creates the given state configuration [9]. The five-dimensional mass chosen as [14] $M_5^2 = \tau(\tau - 4)$, fixes the UV limit of the dual string amplitude with the twist dimension.

In [4], in the context of higher spins mesonic states we showed that the leading infrared contribution for the metric as

$$A(z) = \text{Log}(z \Lambda_{QCD}) + \frac{1 + \sqrt{3}}{2S + \sqrt{3} - 1} \frac{(z \Lambda_{QCD})^2}{1 + e^{(1-z\Lambda_{QCD})}} \quad (5)$$




FIG. 1: Regge trajectory for f_0 (left panel) and pion (right panel) from the Dynamical AdS/QCD model with $\Lambda_{QCD} = 0.3$ GeV (dashed line). Experimental data from [1].

is

$$\mathcal{V}_S(z) \xrightarrow{z \rightarrow \infty} \Lambda_{QCD}^4 (1 + \sqrt{3})^2 z^2, \quad (6)$$

which leads to a satisfactory description of the meson mass spectrum with nearly universal Regge slopes, just using the natural scale of Λ_{QCD} , without any further tuning of parameters. (The spin dependent factor in Eq. (5) is required by universality of effective IR potential (6)). A good analytical approximation to the spectrum calculated with $\Lambda_{QCD} = 0.3$ GeV for $S \geq 1$ and $n \geq 0$ is $m^2 \simeq \frac{1}{10}(11n + 9S + 2)$, that provides an overall fit to the radial and higher-spin excitations [4].

In this letter we propose a generalization of the Dynamical AdS/QCD model for scalar mesons based on the universality of the form of the effective potential in the IR limit. The idea is to make a scale transformation in the holographic distance tailored to fit the slope of the scalar Regge trajectory. We assume the same universal form of the metric as given by

$$A(z) = \text{Log}(z \Lambda_{QCD}) + \frac{(\xi z \Lambda_{QCD})^2}{1 + e^{(1-\xi z \Lambda_{QCD})}} \quad (7)$$

with $\xi = 0.58$ from the fit (see figure 1). The slope of the Regge trajectory is decreased for the lower scalar excitations (see figure 1) in respect to $S = 1$ as $\xi < 1$ and also the ground state $f_0(600)$ is large in the holographic coordinate compared to other light mesons. Therefore, $f_0(600)$ comes as a broad resonance in pionic channels owing to a large overlap between the corresponding amplitudes. More on that will come on what follows (see figure 2 and Table I).

Solving the Sturm-Liouville equation (3) we obtain the spectrum and the wave functions of the scalar mesons states. We used the lowest conformal dimension operator of a scalar, twist 2 [15]. The trajectories found were in agreement of experimental data of f_0 family as shown in figure 1. In this picture the sigma meson is the fundamental state and the other f_0 's are radial excitation of the sigma. We have included in the plot all f_0 's present in the Particle Listing of PDG [1]. That agreement may be




FIG. 2: Normalized wave functions for f_0 mesons (thick solid line) and pion (solid line) from the Dynamical AdS/QCD model with $\Lambda_{\text{QCD}} = 0.3$ GeV.

fortuitous and to support our interpretation we test the physics brought by the structure of these mesons measured by the $S \rightarrow PP$ partial width, that reflects the overlap between the dual string amplitudes of the corresponding mesons. Therefore, the model should include a description of the string dual to the pion state.

The first striking point is the almost vanishing pion mass, and the slope of about 1 GeV^2 for its Regge trajectory, with a value twice of the one found for the scalars. This indicates that the scaling factor of the holographic coordinate for the pseudoscalars should be changed in respect to the f_0 family. This is not a surprise as the pion is a Goldstone boson. To distinguish the pion states in our model, the fifth dimensional mass was rescaled according to $M_5^2 \rightarrow M_5^2 + \lambda z^2$ (see [10] and [15]) and as well the factor ξ in the metric, Eq. (7). The constraints are the pion mass, the slope of the Regge trajectory and the twist 2 from the operator $\bar{q}\gamma^5 q$. The results for the pion Regge trajectory are shown in figure 1 for $\xi = 0.88$ and $\lambda = -2.19 \text{ GeV}^2$. The rescale of the string mass can be an indication of the coupling with the higher twist operators, that in a couple channel model decreases the mass of the ground state. The size of the pion wave function is considerably smaller than the size of sigma as shown in figure 2 which reflects the larger slope for the pseudoscalar Regge trajectory in comparison to the scalar one.

The f_0 's partial decay width into $\pi\pi$ are calculated from the overlap integral of the normalized string amplitudes (Sturm-Liouville form) in the holographic coordinate dual to the scalars (ψ_n) and pion (ψ_π) states,

$$h_n = k \int_0^\infty dz \psi_\pi^2(z) \psi_n(z), \quad (8)$$

where k is a constant with dimension $\sqrt{\text{mass}}$ fitted to the experimental value of the $f_0(1500) \rightarrow \pi\pi$ partial decay

width. The overlap integral is the dual representation of the transition amplitude $S \rightarrow PP$ and therefore the decay width is given by $\Gamma_{\pi\pi}^n = \frac{1}{8\pi} |h_n|^2 \frac{p_\pi}{m_n^2}$, where p_π is the pion momentum in the meson rest frame. The Sturm-Liouville amplitudes of the scalar (pseudoscalar) modes are normalized just as a bound state wave function in quantum mechanics [16, 17], which also corresponds to a normalization of the string amplitude

$$\int_0^\infty dz \psi_m(z) \psi_n(z) = \int_0^\infty dz \varphi_m(z) e^{-(\Phi+3A)} \varphi_n(z) = \delta_{mn}. \quad (9)$$

The overlap integral for the transition amplitude (8) is naturally damped by increasing the f_0 excitation as destructive interference comes into scene within the holographic view. A qualitative understanding of that effect can be seen in figure 2, by observing that within the range of the pion wave function nodes of the higher scalar excitation takes place.

The two-pion partial decay width for the f_0 's present in the particle listing of PDG, are calculated with Eq. (8) and shown in Table I. The width of $f_0(1500)$ is used as normalization. In particular for $f_0(600)$ the model gives a width of about 500 MeV, while its mass is 860 MeV. The range of experimental values quoted in PDG for the sigma mass and width are quite large as depicted in Table I. A recent analysis of the sigma pole in the $\pi\pi$ scattering amplitude from ref.[18] gives $m_\sigma = 441_{-8}^{+16}$ MeV and $\Gamma_\sigma = 544_{-25}^{+18}$ MeV, which in comparison to our results the width seems consistent while the model mass appears somewhat larger. The analysis of the E791 experiment gives $m_\sigma = 478_{-23}^{+24} \pm 17$ MeV and $\Gamma_\sigma = 324_{-40}^{+42} \pm 21$ MeV [19], and the CLEO collaboration [20] quotes $m_\sigma = 513 \pm 32$ MeV and $\Gamma_\sigma = 335 \pm 67$ MeV, both values of the width smaller than our result. Other analysis of the σ -pole in the $\pi\pi \rightarrow \pi\pi$ scattering amplitude present in the decay of heavy mesons indicates a mass around 500 MeV [21]. A rescaling of the string mass as seen necessary for the pion case can lower the sigma mass. We just observe that coupling with higher twist string duals can be a source for the effective decrease of the string mass as in the pion case. The width is not strongly affected as the shift in the string mass mainly dislocates the squared meson mass by a constant. We do not attempt to fine tuning the model at this stage.

The $f_0(980)$ is identified with the first excitation of the string model dual to $q\bar{q}$ state (see Table I). The model mass is shifted to a value above the experimental one, i.e., 1.1 GeV compared to 0.98 GeV. The shift of -0.12 GeV can be attributed to a rescaling of the string mass as in the sigma case. By increasing the excitation of the scalar meson this shift tends to decrease (see $f_0(1500)$ in Table I). The experimental values of $\Gamma_{\pi\pi}$ for $f_0(980)$ is too small compared to our result. We introduced a mixing angle for $f_0(980)$ of $\pm 20^\circ$, that corresponds to a composite nature by mixing, e.g., $s\bar{s}$ with light non-strange quarks[6]. The

mixing angle absolute value between $\sim 12^\circ$ to 28° fits $\Gamma^{\pi\pi}$ within the experimental range.

TABLE I: Two-pion decay width and masses for the f_0 family. Experimental values from PDG[1]. ([†]Mixing angle of 20° . *Fitted value.)

Meson	$M_{exp}(\text{GeV})$	$M_{th}(\text{GeV})$	$\Gamma_{\pi\pi}^{exp}(\text{MeV})$	$\Gamma_{\pi\pi}^{th}(\text{MeV})$
$f_0(600)$	0.4 - 1.2	0.86	600 - 1000	535
$f_0(980)$	0.98 ± 0.01	1.10	$\sim 15 - 80$	$42^\dagger$
$f_0(1370)$	1.2 - 1.5	1.32	$\sim 41 - 141$	141
$f_0(1500)$	1.505 ± 0.006	1.52	38 ± 3	38^*
$f_0(1710)$	1.720 ± 0.006	1.70	$\sim 0 - 6$	5
$f_0(2020)$	1.992 ± 0.016	1.88	—	0.0
$f_0(2100)$	2.103 ± 0.008	2.04	—	1.2
$f_0(2200)$	2.189 ± 0.013	2.19	—	2.5
$f_0(2330)$	2.29 - 2.35	2.33	—	2.8

The mass of the higher scalar excitations are consistent with the experimental data in Table I and figure 1. The two-pion partial decay widths of $f_0(1370)$ and $f_0(1710)$ are compared to the experimental values, and particularly for $f_0(1370)$ we get a result of 141MeV within the experimental range, without any further assumption.

In summary, we provide the basic framework to study the f_0 family given by excitations of the sigma meson as the string duals to $q\bar{q}$ states. The classification, spectroscopy and decay is guided by an Holographic view of the scalar string modes obtained from a Dynamical AdS/QCD model[4] of coupled dilaton-gravity background solution of the Einstein equations. The deformation of the anti-de Sitter metric encodes confinement by the area law behavior of the Wilson loop. Assuming the universality of the metric, apart a scale deformation, for the scalar, pseudoscalar and higher spin string modes dual to $q\bar{q}$ states, the Regge-like spectrum of the radial excitations of f_0 was obtained and the slope about 0.5 GeV^2 fitted to the data. The pion trajectory was also described adding a rescaled string mass. The decrease of the string mass can also improve the description of the lower excitations of the f_0 family, which can be attributed to the coupling of $q\bar{q}$ states to more complex configurations. A nice point about the interpretation of holographic QCD comes from a recent work of Brodsky and Téramond [15], that related the wave equation in the Sturm-Liouville form to the squared mass operator eigenvalue equation for the valence component of the meson light-front wave function. In that respect, the effective potential that we have derived incorporates the coupling of the valence wave function to all other higher Fock-components of the wave function, indicating a pos-

sible new class of AdS/QCD models with the coupling between twist 2 and higher twist modes. The experimentally known partial decay width into $\pi\pi$ give further support to the proposed classification scheme for $f_0(600)$, $f_0(980)$, $f_0(1370)$, $f_0(1500)$ and $f_0(1710)$, while in the particular case of $f_0(980)$ it should mix with more exotic structures as $s\bar{s}$ with an estimated mixing angle around $\pm 20^\circ$.

We acknowledge partial support from the Brazilian agencies CAPES, CNPq and FAPESP.

-
- [1] C. Amsler et al. (Particle Data Group), Phys. Lett. B **667**, 1 (2008) .
 - [2] R. Jaffe, Phys. Rev. D **15**, 28 (1977).
 - [3] L. Maiani, F. Piccinini, A. D. Polosa, V. Riquer, Phys. Rev. Lett. **93**, 212002 (2004); G. 't Hooft, G. Isidori, L. Maiani, A. D. Polosa, V. Riquer, Phys. Lett. B **662**, 424 (2008).
 - [4] W. de Paula, T. Frederico, H. Forkel and M. Beyer, Phys. Rev. D **79**, 075019 (2009), PoS LC2008:046 (2008).
 - [5] U. Gürsoy, E. Kiritsis and F. Nitti, JHEP **0802**, 019 (2008).
 - [6] I. Bediaga, F. Navarra and M. Nielsen, Phys. Lett. B **579**, (2004).
 - [7] G. 't Hooft, Nuclear Physics B **72**, 461 (1974).
 - [8] J. Maldacena, Adv. Theor. Math. Phys. **2**, 231 (1998).
 - [9] J. Polchinski and M. Strassler, Phys. Rev. Lett. **88**, 031601 (2002); H. Boschi, N. Braga and H. Carrion, Eur. Phys. J. C **32**, 529 (2004); Phys. Rev. D **73**, 047901 (2006); G. F. de Téramond and S. J. Brodsky, Phys. Rev. Lett. **94**, 0201601 (2005);
 - [10] H. Forkel, M. Beyer and T. Frederico, JHEP **0707**, 077 (2007); Intl. J. Mod. Phys. E **16**, 2794 (2007).
 - [11] A. Karch, E. Katz, D.T. Son and M.A. Stephanov, Phys. Rev. D **74**, 015005 (2006).
 - [12] A. Vega and I. Schmidt, Phys. Rev. D **78**, 017703 (2008).
 - [13] P. Colangelo, F. De Fazio, F. Giannuzzi, F. Jugeau, and S. Nicotri, Phys. Rev. D **78**, 055009 (2008).
 - [14] E. Witten, Adv. Theor. Math. Phys. **2**, 505 (1998).
 - [15] G. F. de Téramond and S. J. Brodsky, Phys. Rev. Lett. **102**, 081601 (2009).
 - [16] H. Grigoryan and A. Radyushkin, Phys. Rev. D **76**, 095007 (2007).
 - [17] S. J. Brodsky and G. F. de Téramond, Phys. Rev. D **78**, 025032 (2008).
 - [18] I. Caprini, G. Colangelo and H. Leutwyler, Phys. Rev. Lett. **96**, 132001 (2006).
 - [19] E. M. Aitala et al. (Fermilab E791 collaboration), Phys. Rev. Lett. **86**, 770 (2001).
 - [20] H. Muramatsu et al. (CLEO collaboration), Phys. Rev. Lett. **89**, 251802 (2002).
 - [21] D. V. Bugg, Eur. Phys. Jour. C **47**, 57 (2006); AIP Conf. Proc. **1030**, 3 (2008).